

# Poster: Crustaceans

## Region

North Coast, Gascoyne Coast, West Coast, South Coast, Indian Ocean Territories

## Summary

These animals are covered with a protective outer shell so are named crustacean, meaning 'hard-shelled'. Crustaceans belong to the group of animals called arthropods, meaning 'joint-legged'.

**CRUSTACEANS ARMOURD INVERTEBRATES**

**CRUSTY CREATURES**  
These animals are covered with a protective outer shell so are named **crustaceans**, meaning 'hard-shelled'. Crustaceans belong to the group of animals called **arthropods**, meaning 'joint-legged'. This group also includes insects, spiders and scorpions. They are among the most widespread and diverse of all animals and occur in virtually all land, marine and freshwater habitats. There are around 50,000 known crustacean species and scientists estimate there are many more yet to be discovered. The most recognised crustaceans include prawns, lobster and crabs but there are many more, less familiar groups such as barnacles, water fleas and sea lice. Crustaceans range in size from the massive Japanese spider crab, which has a leg span of close to 4 m, to tiny organisms that drift about as part of the plankton.

**CRUSTACEAN CHARACTERISTICS**  
Crustaceans vary a great deal in shape and form, making it very difficult to recognise some of them. In general, however, crustaceans have a hard, external skeleton, which can be thick and tough as with lobsters or thin and transparent as in water fleas. Their limbs are also jointed and can move in nearly every direction. Many also have specialised appendages for feeding, swimming, marking, respiration and reproduction. They have a segmented body – typically divided into a head, thorax (middle segment) and abdomen (tail region). In the majority of larger crustaceans, the head and thorax are fused into a 'cephalothorax' and covered by a single shield-like plate called a carapace. At some stages of their life cycle, crustaceans have two pairs of antennae – unlike other arthropods, which only have one pair. In Hollywood, crustaceans have been used as inspiration for colony computers and robots in some well-known science fiction and movie films.

**BREATHING IN AND OUT OF THE WATER**  
Many crabs can breathe both in and out of the water. In order to breathe in water, they must be kept moist in order to breathe. Respiration in crustaceans can differ. Many of the smaller crustaceans have no special respiratory organs and so gas is exchanged directly through their body surfaces. Larger crustaceans have gills that may be within their bodies or in specialised superlimbs. Some crustaceans can spend long periods out of the water. They keep their gills moist in special gill chambers and when they return from the air, sea monkeys are a type of crustacean called Anomala or lime shrimp. They (along with some hyper-saline larvae and their egg-like cysts) can lay dormant for several years.

**DECAPOD OR NOT**  
The most well-known crustaceans such as crabs, crayfish, lobsters, prawns and shrimps belong to a group called 'decapods', meaning ten legs. Decapods are the largest and most advanced crustaceans, and support several fisheries in Western Australia. Less familiar groups are the non-decapod crustaceans – those that do not have ten legs. These include barnacles, sea lice, lime shrimps, marine shrimps and water fleas. By far the most numerous non-decapod crustaceans are the tiny planktonic animals such as copepods and the larvae of larger crustaceans.

**HOW DO THEY GROW?**  
The hard shell or external skeleton of crustaceans serves as a sort of armour and helps protect them from predators. It also prevents them from growing too other animals. Instead they must periodically shed this exoskeleton in order to increase in size – a process called 'moulting'. A soft new skeleton grows under the old one. When the old skeleton is discarded, it leaves the animal without its main means of protection until the new shell hardens. Many animals are capable of producing four replacement exoskeleton shells, whereas others produce three during their life span.

**DIVERSE DIETS**  
Some crustaceans are filter feeders, collecting tiny plankton and other organic particles of food from the water. Others feed on vegetation or are active predators, seeking out small shellfish and other animals. Many crustaceans are scavengers, feeding on 'detritus' (dead and dying marine life). Several species are even parasitic, often during their larval stages. Crustaceans, in turn, are eaten by many animals, including human beings.

**DID YOU KNOW?**  
It is estimated that 120 million Christmas Island red crabs migrate the island – that equates to nearly a million crabs per square kilometre.  
The Caribbean Periodic cicada is 'Tribolium', Tribolium was one of the first crustaceans, although they were very common. They became extinct 210 million years ago.  
The larvae of many crustaceans are capable of forming hundreds of protective cocoons. Marine shrimp appendages are so strong they can break the spine of an opponent or split a human thumb to the bone with one strike.

**POPULAR CRUSTACEANS**

**BARNACLES**  
Barnacles are hermaphrodites – they have both male and female sexual organs. Barnacles have free-swimming larvae that attach themselves head-down onto hard surfaces including rocks, jetties, boat hulls and even other marine invertebrates. As they grow, their external shell becomes a series of plates, and their jointed legs develop into feathery appendages to feed on plankton.

**LOBSTERS AND CRAYS**  
South-west WA has several species of spiny burrowing crayfish of the European species living within freshwater swamps and seagrasses – and from all these are all listed as 'conservation'. Around eight species of rock lobster are found in Western Australia's tropical and temperate waters, but the most abundant by far is the western rock lobster. This highly crustacean supports one of Western Australia's largest and most valuable fisheries. In WA, the term crayfish is usually given to freshwater species such as murray, gilgus, lacepedes and the introduced jabby.

**CRABS**  
Crabs come in many colours, shapes and sizes and are found in just about every possible habitat, from the intertidal zone to the deep sea – and beneath land! species occur on land. Of the hundreds of species of crabs in WA, the best known is probably the blue swimmer crab, which is caught by the local fish. Others include rock crabs, decapod crabs – which specialise in camouflage, spider crabs and hermit crabs – which inhabit empty mollusk shells.

**PRAWNS AND SHRIMPS**  
Clearer shrimp wait at rocky shores for fish to visit, where they remove and eat debris from the fish. Found in both marine and fresh water, these long-tailed decapods vary in colour – though their shells usually have regular patterns. In Australia, large shrimps are generally known as prawns. Various prawn species, mostly tiger, king and barren prawns, are fished along WA's north-west coast.

**KRILL**  
Krill is a general term used to describe over 80 different species of marine crustaceans known as euphausiids. These small shrimp-like creatures are a vital link in the marine food web between the tiny phytoplankton and larger carnivores such as fish, whales, seals, squid, penguins and many other animals. The most successful animal on the planet in terms of biomass is the Antarctic krill (*Euphausia superba*) – which has a biomass of over 500 million tonnes, or roughly twice that of humans.

**IMPORTANCE**  
The larger decapod crustaceans (such as crabs, lobsters and prawns) are economically valuable. Western Australia has large commercial fisheries of western rock lobster, blue swimmer crab and a number of different prawn species. Blue rock lobster, blue swimmer crab and a number of different prawn species are also popular with recreational fishers. Crustaceans, particularly the small planktonic crustaceans such as krill and copepods, play a major part in marine food webs. Filter-feeding crustaceans eat decomposing plants and animals and act as an important part of the nutrient cycle, particularly in Western Australia.

**REPRODUCTION**  
Many crustaceans exhibit elaborate courtship behaviour, and the males may fight for the chance to mate. In most crustaceans, the oviduct is separate, with females producing eggs and males producing sperm. In many species, females brood the eggs beneath their abdomen (that is, they carry the eggs until they hatch). These females are said to be 'broody' when carrying eggs. After fertilisation the eggs can take anywhere from several days to several weeks to develop and hatch, depending on the species. Some crustaceans hatch young that immediately look like smaller versions of the adult. Most however, go through a free-swimming larval stage called a 'nauplius'. After a series of changes (called 'metamorphosis'), the larvae eventually look like a miniature adult. They continue to feed and grow into 'juvveniles' (adult animals that are not yet able to reproduce).

## Download Resource

[Crustacean Poster](#)